


Gibbs Energy of Formation of Ammonia

Here, we calculate the Gibbs energy of formation of ammonia (NH₃), given the reaction



Values of the Gibbs energy of formation tabulated in the literature are normally only given at standard temperature, or for a small number of temperatures. However, the [ThermophysicalData:-Chemicals](#) package contains curve fits for experimental values of enthalpy and entropy of N₂, H₂ and NH₃, correlated against temperature.

This means that Maple will help you calculate the Gibbs energy of formation of ammonia at any temperature (assuming that the enthalpy and entropy are within the bounds of the fitted data).

> *restart* :
with(*ThermophysicalData:-Chemicals*) :

Temperature

> $T := 298.15 \text{ K}$:

Enthalpy

> $h_{\text{N2}} := \text{Property}(\text{"Hmolar"}, \text{"N2"}, \text{"temperature"} = T);$
 $h_{\text{H2}} := \text{Property}(\text{"Hmolar"}, \text{"H2"}, \text{"temperature"} = T);$
 $h_{\text{NH3}} := \text{Property}(\text{"Hmolar"}, \text{"NH3"}, \text{"temperature"} = T);$

$$9.92 \times 10^{-6} \frac{\text{J}}{\text{mol}}$$

$$-4.96 \times 10^{-6} \frac{\text{J}}{\text{mol}}$$

$$-45.94 \times 10^3 \frac{\text{J}}{\text{mol}}$$

(1)

Entropy

> $s_{\text{N2}} := \text{Property}(\text{"Smolar"}, \text{"N2"}, \text{"temperature"} = T);$
 $s_{\text{H2}} := \text{Property}(\text{"Smolar"}, \text{"H2"}, \text{"temperature"} = T);$
 $s_{\text{NH3}} := \text{Property}(\text{"Smolar"}, \text{"NH3"}, \text{"temperature"} = T);$

$$\begin{aligned}
 &191.61 \frac{\text{J}}{\text{mol K}} \\
 &130.68 \frac{\text{J}}{\text{mol K}} \\
 &192.77 \frac{\text{J}}{\text{mol K}}
 \end{aligned}
 \tag{2}$$

Change in enthalpy and entropy per mole of NH_3

$$\begin{aligned}
 &> \Delta H := 0.5 (2 h_{\text{NH}_3} - (h_{\text{N}_2} + 3 h_{\text{H}_2})); \\
 &\Delta S := 0.5 (2 s_{\text{NH}_3} - (s_{\text{N}_2} + 3 s_{\text{H}_2})); \\
 &\quad -45.94 \frac{\text{kJ}}{\text{mol}} \\
 &\quad -99.06 \frac{\text{J}}{\text{mol K}}
 \end{aligned}
 \tag{3}$$

Hence the Gibbs Energy of Formation

$$\begin{aligned}
 &> \Delta G := \Delta H - T \cdot \Delta S \\
 &\quad -16.41 \frac{\text{kJ}}{\text{mol}}
 \end{aligned}
 \tag{4}$$

Legal Notice: © Maplesoft, a division of Waterloo Maple Inc. 2018. Maplesoft and Maple are trademarks of Waterloo Maple Inc. This application may contain errors and Maplesoft is not liable for any damages resulting from the use of this material. This application is intended for non-commercial, non-profit use only. Contact Maplesoft for permission if you wish to use this application in for-profit activities.